

**AUTHORS INDEX of QIC Vol.13 (2013)**

<b>A-H</b>	<b>H-N</b>	<b>N-Z</b>
<p>J. Aamlof, <i>Fidelity as a figure of merit in quantum error correction</i> (1&amp;2) 9</p> <p>A. Abdollahi, <i>Reversible logic synthesis by quantum rotation gates</i> (9&amp;10) 771</p> <p>A. Acin, see G. Prettico</p> <p>A. Ahanj, see S.Y. Mirafzali</p> <p>B. Amento (I), <i>Quantum binary field inversion: improved circuit depth via choice of basis representation</i> (1&amp;2) 116</p> <p>B. Amento (II), <i>Efficient quantum circuits for binary elliptic curve arithmetic: reducing T-gate complexity</i> (7&amp;8) 631</p> <p>A. Anshu, <i>Pseudo-telepathy games using graph states</i> (9&amp;10) 833</p> <p>D.M. Appleby, <i>Galois automorphisms of a symmetric measurement</i> (7&amp;8) 672</p> <p>N. de Beaudrap, <i>A linearized stabilizer formalism for systems of finite dimension</i> (1&amp;2) 73</p> <p>E.C. Behrman, <i>Multiqubit entanglement of a general input state</i> (1&amp;2) 36</p> <p>G. Bjork, see J. Aamlof</p> <p>F.G.S.L. Brandao (I), see L. Pankowski</p> <p>F.G.S.L. Brandao (II), <i>Exponential quantum speed-ups are generic</i> (11&amp;12) 901</p> <p>S. Bravyi, <i>Subsystem surface codes with three-qubit check operators</i> (11&amp;12) 963</p> <p>J. Briet, <i>Multipartite entanglement in XOR games</i> (3&amp;4) 334</p> <p>J. Brown, <i>Perfect state transfer on signed graphs</i> (5&amp;6) 511</p> <p>H. Buhrman, see J. Briet</p> <p>A. Chailloux, <i>Lower bounds for quantum oblivious transfer</i> (1&amp;2) 158</p> <p>B. Chen, <i>A note on locally unextendible non-maximally entangled basis</i> (11&amp;12) 1077</p> <p>J.-L. Chen, see C.-Q. Pang</p> <p>R.-X. Chen, see S.-B. Zheng (II)</p> <p>W. Chen, see Z.-Q. Yin</p> <p>X.Y. Chen, see L.T. Shen</p> <p>N. Delfosse, <i>Upper bounds on the rate of low density stabilizer codes for the quantum erasure channel</i> (9&amp;10) 793</p> <p>M. Demianowicz, <i>Multiaccess quantum communication and product higher rank numerical range</i> (7&amp;8) 541</p> <p>G. Duclos-Cianci, see S. Bravyi</p> <p>W. Dur, see M. Skotiniotis</p> <p>B. Eastin, see A.M. Meier</p> <p>C. Eltschka, <i>Optimal class-specific witnesses for three-qubit</i></p>	<p>M.B. Hastings (II), <i>Obstructions to classically simulating the quantum adiabatic algorithm</i> (11&amp;12) 1038</p> <p>Y. He, <i>Optical detection of quantum entanglement between two quantum dots near a metal nanoparticle</i> (3&amp;4) 324</p> <p>M. Hellmund, <i>Entanglement and output entropy of the diagonal map</i> (5&amp;6) 379</p> <p>B. Helou, see T. Jochym-O'Connor</p> <p>D.A. Herrera-Marti, <i>Loss tolerance with a concatenated graph state</i> (11&amp;12) 995</p> <p>C.D. Hill, <i>Fault-tolerant quantum error correction code conversion</i> (5&amp;6) 439</p> <p>L.C.L. Hollenberg, see C.D. Hill</p> <p>K. Horodecki, see P. Joshi</p> <p>M. Horodecki (I), see P. Joshi</p> <p>M. Horodecki (II), see L. Pankowski</p> <p>M. Horodecki (III), see F.G.S.L. Brandao (II)</p> <p>P. Horodecki (I), see M. Demianowicz</p> <p>P. Horodecki (II), see P. Joshi</p> <p>R. Horodecki, see P. Joshi</p> <p>Y. Huang, see C.-H. Fan</p> <p>T. Ito, see D. Gavinsky</p> <p>K. Javidan, see S.Y. Mirafzali</p> <p>C.-G. Ji, <i>Spin squeezing of one-axis twisting model in the presence of phase dephasing</i> (3&amp;4) 266</p> <p>Y. Jiang, see C.-Q. Pang</p> <p>G.-R. Jin, see C.-G. Ji</p> <p>T. Jochym-O'Connor, <i>The robustness of magic state distillation against errors in Clifford gates</i> (5&amp;6) 361</p> <p>P. Joshi, <i>No-broadcasting of non-signalling boxes via operations which transform local boxes into local ones</i> (7&amp;8) 567</p> <p>D. Kaszlikowski, see A. Kay</p> <p>A. Kay, <i>Optimal asymmetric quantum cloning for quantum information and computation</i> (9&amp;10) 880</p> <p>I. Kerenidis, see A. Chailloux</p> <p>V. Kliuchnikov, <i>Fast and efficient exact synthesis of single-qubit unitaries generated by Clifford and T gates</i> (7&amp;8) 607</p> <p>E. Knill, see A.M. Meier</p> <p>B. Kraus, see M. Skotiniotis</p> <p>R. Laflamme, see T. Jochym-O'Connor</p> <p>R. Li, <i>Hermitian dual containing BCH codes and Construction of new quantum codes</i> (1&amp;2) 21</p> <p>M.-L. Liang, see C.-Q. Pang</p> <p>T. Lee, see J. Briet</p> <p>H.-W. Li, see Z.-Q. Yin</p> <p>E.H. Lieb, <i>Upper bounds on mixing rates</i> (11&amp;12) 986</p> <p>Y. Liu, see R. Li</p>	<p>X. Ni, <i>Commuting quantum circuits: efficiently classical simulations versus hardness results</i> (1&amp;2) 54</p> <p>H. Nizamidin, see B. Chen</p> <p>C.-Q. Pang, <i>Most robust and fragile two-qubit entangled states under depolarizing channels</i> (7&amp;8) 645</p> <p>L. Pankowski, <i>Entanglement distillation by extendible maps</i> (9&amp;10) 751</p> <p>M. Pedram, see A. Abdollahi</p> <p>P. Pham, <i>A 2D nearest-neighbor quantum architecture for factoring in polylogarithmic depth</i> (11&amp;12) 937</p> <p>S.J.D. Phoenix, see F. Shah Khan</p> <p>D. Poulin, see S. Bravyi</p> <p>J. Preskill (I), <i>Sufficient condition on noise correlations for scalable quantum computing</i> (3&amp;4) 181</p> <p>J. Preskill (II), see J. Napp</p> <p>G. Prettico, <i>Can bipartite classical information resources be activated?</i> (3&amp;4) 245</p> <p>R. Ramanathan, see A. Kay</p> <p>A. Raz, see J. Brown</p> <p>X.-J. Ren, <i>Non-monogamy of quantum discord and upper bounds for quantum correlation</i> (5&amp;6) 469</p> <p>M. Rötteler (I), see B. Amento (I)</p> <p>M. Rötteler (II), see B. Amento (II)</p> <p>T. Rudolph, see D.A. Herrera-Marti</p> <p>M. Saeedi, see A. Abdollahi</p> <p>P.J. Salas, <i>Security of plug-and-play QKD arrangements with finite resources</i> (9&amp;10) 861</p> <p>M. Sarbisei, see S.Y. Mirafzali</p> <p>I. Sargolzahi, see S.Y. Mirafzali</p> <p>F. Shah Khan, <i>Gaming the quantum</i> (3&amp;4) 231</p> <p>K. Scharnhorst, <i>Entanglement capabilities of the spin representation of (3+1)D-conformal transformations</i> (11&amp;12) 925</p> <p>L.T. Shen, <i>Cooling distant atoms into steady entanglement via coupled cavities</i> (3&amp;4) 281</p> <p>J. Siewert, see C. Eltschka</p> <p>J. Sikora (I), see S. Gharibian</p> <p>J. Sikora (II), see A. Chailloux</p> <p>M. Skotiniotis, <i>Efficient quantum communication under collective noise</i> (3&amp;4) 290</p> <p>G. Smith, see L. Pankowski</p> <p>J.E. Steck, see E.C. Behrman</p> <p>R. Steinwandt (I), see B. Amento (I)</p> <p>R. Steinwandt (II), see B. Amento (II)</p> <p>M. Suchara, see S. Bravyi</p> <p>Z. Sun, see C.-H. Fan</p> <p>K.M. Svore, see P. Pham</p> <p>C. Tamon, see J. Brown</p> <p>S. Upadhyay, see S. Gharibian</p> <p>A. Vershynina, see E.H. Lieb</p> <p>T. Vidick, see J. Briet</p>

<p><i>entanglement from Greenberger-Horne-Zeilinger symmetry</i> (3&amp;4) 210</p> <p>C.-H. Fan, <i>Quantum discord and quantum phase transition in spin-1/2 frustrated Heisenberg chain</i> (5&amp;6) 452</p> <p>H. Fan, see X.-J. Ren</p> <p>S.-M. Fei, see B. Chen</p> <p>S.T. Flammia, <i>Counterexamples to Kalai's conjecture C</i> (1&amp;2) 1</p> <p>A.G. Fowler, see C.D. Hill</p> <p>D. Gavinsky, <i>Quantum fingerprints that keep secrets</i> (7&amp;8) 583</p> <p>M. Gharibi, <i>Reduction from non-injective hidden shift problem to injective hidden shift problem</i> (3&amp;4) 221</p> <p>S. Gharibian, <i>QMA variants with polynomially many provers</i> (1&amp;2) 135</p> <p>C. Godsil, see J. Brown</p> <p>G.-C. Guo, see Z.-Q. Yin</p> <p>A. Grudka, see P. Joshi</p> <p>S. Hallgren, <i>The local Hamiltonian problem on a line with eight states is QMA-complete</i> (9&amp;10) 721</p> <p>Z.-F. Han, see Z.-Q. Yin</p> <p>A.W. Harrow, see S.T. Flammia</p> <p>M. Hastings (I), <i>Trivial low energy states for commuting Hamiltonians, and the quantum PCP conjecture</i> (5&amp;6) 393</p>	<p>Y.-C. Liu, see C.-G. Ji</p> <p>T. Machida (I), <i>Realization of the probability laws in the quantum central limit theorems by a quantum walk</i> (5&amp;6) 430</p> <p>T. Machida (II), <i>Limit theorems for the interference terms of discrete-time quantum walks on the line</i> (7&amp;8) 661</p> <p>S. Maitra, <i>Cryptanalysis of a secret sharing scheme</i> (1&amp;2) 178</p> <p>D. Mallory, see J. Brown</p> <p>D. Maslov, see V. Kliuchnikov</p> <p>A.M. Meier, <i>Magic-state distillation with the four-qubit code</i> (3&amp;4) 195</p> <p>M. Mhalla, see A. Anshu</p> <p>S.Y. Mirafzali, <i>Measurement-induced nonlocality for an arbitrary bipartite state</i> (5&amp;6) 479</p> <p>M. Mosca, see V. Kliuchnikov</p> <p>D. Nagaj, see S. Hallgren</p> <p>K. Nandi, see S. Maitra</p> <p>J. Napp, <i>Optimal Bacon-Shor codes</i> (5&amp;6) 490</p> <p>S. Narayanaswami, see S. Hallgren</p> <p>M. Van den Nest (I), see X. Ni</p> <p>M. Van den Nest (II), <i>Efficient classical simulations of quantum Fourier transforms and Normalizer circuits over Abelian groups</i> (11&amp;12) 1007</p>	<p>D.S. Wang, see C.D. Hill</p> <p>S. Wang, see Z.-Q. Yin</p> <p>Z. Wei, <i>Full characterization of quantum correlated equilibria</i> (9&amp;10) 846</p> <p>H.-Z. Wu, see L.T. Shen</p> <p>H.-N. Xiong, see C.-H. Fan</p> <p>Z. Xu, see R. Li</p> <p>H. Yadsan-Appleby, see D.M. Appleby</p> <p>Z.-B. Yang, see L.T. Shen</p> <p>Z.-Q. Yin, <i>A study of BB84 protocol in a device-independent scenario: from the view of entanglement distillation</i> (9&amp;10) 827</p> <p>Y. Yu, see T. Jochym-O'Connor</p> <p>G. Zauner, see D.M. Appleby</p> <p>G. Zemor, see N. Delfosse</p> <p>F.-L. Zhang, see C.-Q. Pang</p> <p>S. Zhang, see Z. Wei</p> <p>S.-B. Zheng (I), see L.T. Shen</p> <p>S.-B. Zheng (II), <i>Entanglement transfer between atomic qubits and thermal fields</i> (5&amp;6) 531</p> <p>K.-D. Zhu, see Y. He</p> <p>F. Zou, see R. Li</p> <p>K. Zyczkowski, see M. Demianowicz</p>
--	---	---

\* in the order: first Author's name, *article title*, (issue no.) starting page number